

MTH5113 (Winter 2022): Problem Sheet 5

All coursework should be submitted individually.

- Problems marked “[**Marked**]” should be submitted and will be marked.

Be sure to write your **name** and **student ID** on your submission!

Please submit the completed problem(s) on QMPlus:

- At the portal for **Coursework Submission 3**.
-

(1) (*Warm-up*) Given a curve C , along with a pair of parametrisations γ_1 and γ_2 of C :

- (i) Compute, for each parameter $\mathbf{t}$, the unit tangent vectors

$$\frac{1}{|\gamma_1'(\mathbf{t})|} \cdot \gamma_1'(\mathbf{t})_{\gamma_1(\mathbf{t})}, \quad \frac{1}{|\gamma_2'(\mathbf{t})|} \cdot \gamma_2'(\mathbf{t})_{\gamma_2(\mathbf{t})}.$$

- (ii) Determine whether γ_1 and γ_2 generate the same orientation or opposite orientations of C . Give a brief justification of your answer.

(a) *Circle*: $C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$, and

$$\begin{aligned} \gamma_1 : \mathbb{R} &\rightarrow C, & \gamma_1(\mathbf{t}) &= (\cos \mathbf{t}, \sin \mathbf{t}), \\ \gamma_2 : \mathbb{R} &\rightarrow C, & \gamma_2(\mathbf{t}) &= (-\sin \mathbf{t}, -\cos \mathbf{t}). \end{aligned}$$

(b) *Line*: $C = \{(x, y, z) \in \mathbb{R}^3 \mid x + y = 2, x + z = 1\}$, and

$$\begin{aligned} \gamma_1 : \mathbb{R} &\rightarrow C, & \gamma_1(\mathbf{t}) &= (\mathbf{t}, 2 - \mathbf{t}, 1 - \mathbf{t}), \\ \gamma_2 : \mathbb{R} &\rightarrow C, & \gamma_2(\mathbf{t}) &= (2 - \mathbf{t}, \mathbf{t}, \mathbf{t} - 1). \end{aligned}$$

(2) (*Warm-up*) Find both the unit tangents and the unit normals to each of the following curves C at the given point $\mathbf{p}$.

(a) $C = \{(t^3, t) \in \mathbb{R}^2 \mid t \in \mathbb{R}\}$ and $\mathbf{p} = (-8, -2)$.

(b) $C = \{(x, y) \in \mathbb{R}^2 \mid x^4 + 2y^2 = 3\}$ and $\mathbf{p} = (-1, 1)$.

(3) (*Warm-up*) Consider the following regular parametric curve:

$$\mathbf{b} : (0, 1) \rightarrow \mathbb{R}^2, \quad \mathbf{b}(t) = \left(t, \frac{2}{3} t^{\frac{3}{2}} \right).$$

(a) Compute the arc length of $\mathbf{b}$.

(b) Compute the curve integral of the function F over $\mathbf{b}$, where

$$F : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad F(x, y) = 1 + x.$$

(c) Compute the curve integral of the function G over $\mathbf{b}$, where

$$G : \mathbb{R} \times (0, \infty) \rightarrow \mathbb{R}, \quad G(x, y) = \frac{2}{3} + \frac{y}{\sqrt{x}}.$$

(4) [Marked] Consider the following ellipse,

$$C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + 4y^2 = 4\},$$

and consider the following function,

$$F : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad F(x, y) = \sqrt{1 + 3y^2}.$$

(a) Give an *injective* parametrisation γ of C such that *the image of γ differs from C by only a finite number of points*. Be sure to specify the domain of γ ! (Hint: Recall all the possible solutions to Question (5) of Problem Sheet 4.)

(b) Compute the curve integral of F over C .

(5) [Tutorial] Let us derive some (possibly) familiar formulas!

(a) Let $\mathcal{C}$ denote the unit circle,

$$\mathcal{C} = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}.$$

Show that at each $\mathbf{p} \in \mathcal{C}$, the unit normals to $\mathcal{C}$ at $\mathbf{p}$ are precisely $\pm \mathbf{p}$.

(b) Let G_f denote the graph of a function $f : (a, b) \rightarrow \mathbb{R}$:

$$G_f = \{(x, y) \in \mathbb{R}^2 \mid y = f(x), a < x < b\}.$$

Find a formula for the arc length of G_f .

(c) Let $\rho : (c, d) \rightarrow \mathbb{R}$, and consider the following *polar parametric curve*:

$$\lambda_\rho : (c, d) \rightarrow \mathbb{R}^2, \quad \lambda_\rho(\theta) = (\rho(\theta) \cos \theta, \rho(\theta) \sin \theta).$$

Find a formula for the arc length of λ_ρ .

(6) (*Issues with parametrisations*) Let H denote the curve

$$H = \{(x, y, z) \in \mathbb{R}^3 \mid x = \cosh z, y = \sinh z\},$$

and let γ be the parametric curve

$$\gamma : \mathbb{R} \rightarrow \mathbb{R}^3, \quad \gamma(t) = (\cosh t, \sinh t, t).$$

(For this problem, you may assume that you already know H is a curve.)

- (a) What statements must you prove in order to show that γ is a parametrisation of H , according to the definition given in this module?
- (b) Show that γ is indeed a parametrisation of H .
- (c) Oh no, Mr. Error (from question (6) of *Problem Sheet 4*) is back to his erroneous ways! He decides to describe the points of H using the parametric curve

$$\zeta : \mathbb{R} \rightarrow H, \quad \zeta(t) = (\cosh t^2, \sinh t^2, t^2).$$

He computes (correctly) that

$$\zeta(0) = (1, 0, 0) \in H, \quad \zeta'(0) = (0, 0, 0).$$

He then (incorrectly) concludes that $T_{(1,0,0)}H$ contains only one element,

$$T_{(1,0,0)}H = T_\zeta(0) = \{s \cdot \zeta'(0)_{\zeta(0)} \mid s \in \mathbb{R}\} = \{(0, 0, 0)_{(1,0,0)}\},$$

hence it is a 0-dimensional space! What error did Mr. Error make this time?

(7) (*Orient my hyperbola!*) Let $\mathcal{H}$ denote the *hyperbola*,

$$\mathcal{H} = \{(x, y) \in \mathbb{R}^2 \mid x^2 - y^2 = 1\}.$$

Describe all the possible orientations of $\mathcal{H}$. How many such orientations are there?

(8) (*Arc length parametrisations*) Let (a, b) be a finite open interval, and let $\gamma : (a, b) \rightarrow \mathbb{R}^n$ be a regular parametric curve. We can then define the change of variables

$$s = \phi(t) = \int_a^t |\gamma'(\tau)| \, d\tau.$$

Note that s represents the total length travelled by γ up to parameter t .

(a) Show that the following holds for any $t \in (a, b)$:

$$\frac{ds}{dt} = |\gamma'(t)|.$$

The reparametrisation λ of γ defined as

$$\lambda : (0, L(\gamma)) \rightarrow \mathbb{R}^n, \quad \lambda(s) = \gamma(t)$$

is called the *arc length reparametrisation*, since its parameter is itself the distance travelled.

(b) Let $R > 0$, and let γ be the regular parametric curve

$$\gamma : (0, 2\pi) \rightarrow \mathbb{R}^2, \quad \gamma(t) = (R \cos t, R \sin t).$$

Find the arc length reparametrisation λ of this γ . What is the domain of λ ?

(9) (*Parental advisory, implicit content*) (*Not examinable*) A special case of the *implicit function theorem* for functions of two variables can be stated as follows:

Theorem. (*Implicit Function Theorem*) Let $U \subseteq \mathbb{R}^2$ be open and connected, let $f : U \rightarrow \mathbb{R}$ be smooth, and let C denote the level set

$$C = \{(x, y) \in U \mid f(x, y) = c\}, \quad c \in \mathbb{R}.$$

Suppose, in addition, that $(x, y) \in C$ and that $\partial_2 f(x, y) \neq 0$. Then, there exists some open set $V \subseteq \mathbb{R}^2$, with $(x, y) \in V$, such that $C \cap V$ is the graph of a function, i.e.,

$$C \cap V = \{(x, h(x)) \mid x \in I\},$$

where I is an open interval, and where $h : I \rightarrow \mathbb{R}$ is smooth.

In addition, an analogous theorem holds with the roles of $\mathbf{x}$ and $\mathbf{y}$ interchanged.

- (a) How does the above implicit function theorem relate to the process of implicit differentiation that you learned in calculus? (An informal description will suffice here.)
- (b) How does the above implicit function theorem relate to the proof of Theorem 3.26 in the lecture notes? Again, an informal description will suffice here.