

MTH5113 (Winter 2022): Problem Sheet 6

All coursework should be submitted individually.

- Problems marked “[**Marked**]” should be submitted and will be marked.

Be sure to write your **name** and **student ID** on your submission!

Please submit the completed problem(s) on QMPlus:

- At the portal for **Coursework Submission 3**.
-

(1) (*Warm-up*) For each of the sets C and points $\mathbf{p} \in C$ given below:

- (i) Show that C is a curve.
- (ii) Sketch C , and indicate the point $\mathbf{p}$ on C .
- (iii) Give a parametrisation of C that passes through $\mathbf{p}$.

(a) *Hyperbola*:

$$C = \{(x, y) \in \mathbb{R}^2 \mid x^2 - y^2 = -1\}, \quad \mathbf{p} = (0, -1).$$

(b) *Cubic*:

$$C = \{(x, y) \in \mathbb{R}^2 \mid (x - 2)^3 = y - 3\}, \quad \mathbf{p} = (0, -5).$$

(c) *Ellipse*:

$$C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + 6x + 4y^2 - 8y = 3\}, \quad \mathbf{p} = (-3, -1).$$

(2) (*Fun with plotting*) The following are exercises involving sketching parametric surfaces. Do make use of computer programs or webpages (see the links in the *Additional Resources* section on the *QMPlus* page) to help you with your sketches.

(a) *Sphere*: Consider the following parametric surface:

$$\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad \sigma(\mathbf{u}, \mathbf{v}) = (\cos \mathbf{u} \sin \mathbf{v}, \sin \mathbf{u} \sin \mathbf{v}, \cos \mathbf{v}).$$

- (i) Sketch the paths obtained from σ by holding $\mathbf{v}$ constant, with values

$$\mathbf{v}_0 = 0, \frac{\pi}{4}, \frac{\pi}{2}, \frac{3\pi}{4}, \pi.$$

(ii) Sketch the paths obtained from σ by holding u constant, with values

$$u_0 = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi.$$

(iii) Sketch the image of σ .

(b) *Hyperboloid*: Consider the following parametric surface:

$$\mathbf{h} : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad \mathbf{h}(u, v) = (\cos u \cosh v, \sin u \cosh v, \sinh v).$$

(i) Sketch the paths obtained from $\mathbf{h}$ by holding v constant, with values

$$v_0 = -2, -1, 0, 1, 2.$$

(ii) Sketch the paths obtained from $\mathbf{h}$ by holding u constant, with values

$$u_0 = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi.$$

(iii) Sketch the image of $\mathbf{h}$.

(3) (*Warm-up*) For each of the following parametric surfaces σ and parameters (u_0, v_0) , compute the tangent plane to σ at (u_0, v_0) :

(a) σ is the parametric *cylinder*,

$$\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad \sigma(u, v) = (\cos u, \sin u, v),$$

$$\text{and } (u_0, v_0) = \left(\frac{\pi}{2}, -1\right).$$

(b) σ is the parametric *one-sheeted hyperboloid*,

$$\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad \sigma(u, v) = (\cos u \cosh v, \sin u \cosh v, \sinh v).$$

$$\text{and } (u_0, v_0) = (\pi, 1).$$

(4) (*Introduction to curve integrals*) One can also define an intermediate notion of curve integration of vector fields over *parametric curves*. More specifically:

Definition. Let $\gamma : (a, b) \rightarrow \mathbb{R}^n$ be a parametric curve, and let $\mathbf{F}$ be a vector field that is defined on the image of γ . We then define the *curve integral* of $\mathbf{F}$ over γ by

$$\int_{\gamma} \mathbf{F} \cdot d\mathbf{s} = \int_a^b [\mathbf{F}(\gamma(t)) \cdot \gamma'(t)_{\gamma(t)}] dt.$$

For each of the following γ and $\mathbf{F}$, compute the curve integral of $\mathbf{F}$ over γ :

(a) γ is the regular parametric curve

$$\gamma : (0, 1) \rightarrow \mathbb{R}^3, \quad \gamma(t) = (t, -t, 2t),$$

and $\mathbf{F}$ is the vector field on $\mathbb{R}^3$ given by

$$\mathbf{F}(x, y, z) = (x, y, z)_{(x, y, z)}.$$

(b) γ is the regular parametric curve

$$\gamma : (0, 2\pi) \rightarrow \mathbb{R}^2, \quad \gamma(t) = (\cos t, \sin t \cos t),$$

and $\mathbf{F}$ is the vector field on $\mathbb{R}^2$ given by

$$\mathbf{F}(x, y) = (x^2, 0)_{(x, y)}.$$

(5) [Marked] Consider the *clockwise-oriented* (shifted) *circle*,

$$C = \{(x, y) \in \mathbb{R}^2 \mid (x + 2)^2 + (y - 3)^2 = 9\},$$

and consider the vector field $\mathbf{F}$ on $\mathbb{R}^2$ given by

$$\mathbf{F}(x, y) = (y, -x)_{(x, y)}.$$

(a) Give an *injective* parametrisation γ of C such that *the image of γ differs from C by only a finite number of points*. Which orientation of C does γ generate?

(b) Compute the (curve) integral of $\mathbf{F}$ over the curve C .

(6) [Tutorial] For each of the following oriented curves C and vector fields $\mathbf{F}$:

- (i) Give an *injective* parametrisation γ of C such that *the image of γ differs from C by only a finite number of points*. Which orientation does γ generate?
- (ii) Compute the (curve) integral of $\mathbf{F}$ over the curve C .

(a) C is the *anticlockwise-oriented ellipse*,

$$C = \{(x, y) \in \mathbb{R}^2 \mid 3x^2 + 2y^2 = 6\},$$

and $\mathbf{F}$ is the vector field on $\mathbb{R}^2$ given by

$$\mathbf{F}(x, y) = (y, -x)_{(x,y)}.$$

(b) C is the *downward-oriented* (with decreasing z -value) *helical segment*,

$$C = \{(\cos t, \sin t, t) \mid t \in (0, 2\pi)\},$$

and $\mathbf{F}$ is the vector field on $\mathbb{R}^3$ given by

$$\mathbf{F}(x, y, z) = (-y, x, 1)_{(x,y,z)}.$$

(7) (*Exploring curvature*) Let $\gamma : I \rightarrow \mathbb{R}^n$ be any regular parametric curve. We then define the *curvature* of γ at $t \in I$ by the formula

$$\kappa_\gamma(t) = \frac{1}{|\gamma'(t)|} \left| \left(\frac{\gamma'}{|\gamma'|} \right)'(t) \right|.$$

(This can be viewed as the “change in the direction of γ per unit length”; see the 2019 version of the *MTH5113 lecture notes* for additional discussions of curvature.)

(a) Let $\mathbf{p}, \mathbf{v} \in \mathbb{R}^n$, and let ℓ be the parametric line,

$$\ell : \mathbb{R} \rightarrow \mathbb{R}^n, \quad \ell(t) = \mathbf{p} + t\mathbf{v}.$$

Compute the curvature of ℓ at every $t \in \mathbb{R}$.

(b) Let $R > 0$, and let γ_R be the parametric circle of radius R :

$$\gamma_R : \mathbb{R} \rightarrow \mathbb{R}^2, \quad \gamma_R(t) = (R \cos t, R \sin t).$$

Compute the curvature of γ_R at every $t \in \mathbb{R}$.

- (c) Show that *curvature is independent of parametrisation*. More specifically, show that if γ is a reparametrisation of $\tilde{\gamma} : \tilde{I} \rightarrow \mathbb{R}^n$, with corresponding change of variables $\phi : I \rightarrow \tilde{I}$ (in particular, $\gamma(t) = \tilde{\gamma}(\phi(t))$ for all $t \in I$), then

$$\kappa_\gamma(t) = \kappa_{\tilde{\gamma}}(\phi(t)), \quad t \in I.$$

- (8) (*Polar curves*) Let $h : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth positive periodic function, with period 2π :

$$h(\theta) > 0, \quad h(\theta + 2\pi) = h(\theta), \quad \theta \in \mathbb{R}.$$

Let the *polar curve* P be the set of all points in $\mathbb{R}^2$ satisfying the relation

$$r = h(\theta)$$

in polar coordinates. (Here, you can assume that P is indeed a curve.)

- (a) The unit circle $\mathcal{C} = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ is a polar curve. What is h here?
- (b) Give an injective parametrisation of P whose image is all of P except for a single point.
- (c) Derive a formula for the arc length of P .

- (9) (*Conic sections*) Let N denote the following cone:

$$N = \{(x, y, z) \in \mathbb{R}^3 \mid z^2 = x^2 + y^2\}.$$

In addition, let $P \subseteq \mathbb{R}^3$ denote an arbitrary plane that does not pass through the origin. More specifically, P is a set of the form

$$P = \{(x, y, z) \in \mathbb{R}^3 \mid ax + by + cz = d\}$$

where $a, b, c, d \in \mathbb{R}$ satisfy $(a, b, c) \neq (0, 0, 0)$ and $d \neq 0$. A set of the form $N \cap P$ (i.e. the intersection of the cone N and the plane P) is called a *conic section*.

- (a) Use the theorem in Question (9) of Problem Sheet 4 to show that any conic section $N \cap P$ is indeed a curve. (*Hint: You will have to be resourceful to do this. The first step is to express $N \cap P$ as an appropriate level set.*)

(b) Find examples of such planes $\mathbf{P}$ such that the conic section $\mathbf{N} \cap \mathbf{P}$ is:

- (i) A circle.
- (ii) An ellipse.
- (iii) A parabola.
- (iv) A hyperbola.

Check your answers graphically on a computer!