

MTH5113 (Winter 2022): Problem Sheet 7

All coursework should be submitted individually.

- Problems marked “[Marked]” should be submitted and will be marked.

Be sure to write your **name** and **student ID** on your submission!

Please submit the completed problem(s) on QMPlus:

- At the portal for **Coursework Submission 4**.
-

(1) (*Warm-up*) For each of the parametric surfaces σ given below and every pair of parameters (\mathbf{u}, \mathbf{v}) in the domain of σ , compute the following:

- (i) $\partial_1 \sigma(\mathbf{u}, \mathbf{v})$ and $\partial_2 \sigma(\mathbf{u}, \mathbf{v})$.
- (ii) $\partial_1 \sigma(\mathbf{u}, \mathbf{v}) \times \partial_2 \sigma(\mathbf{u}, \mathbf{v})$.
- (iii) $|\partial_1 \sigma(\mathbf{u}, \mathbf{v}) \times \partial_2 \sigma(\mathbf{u}, \mathbf{v})|$.
- (a) *Sphere*: $\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, where $\sigma(\mathbf{u}, \mathbf{v}) = (\cos \mathbf{u} \sin \mathbf{v}, \sin \mathbf{u} \sin \mathbf{v}, \cos \mathbf{v})$.
- (b) *Torus*: $\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3$, where $\sigma(\mathbf{u}, \mathbf{v}) = ((2 + \cos \mathbf{u}) \cos \mathbf{v}, (2 + \cos \mathbf{u}) \sin \mathbf{v}, \sin \mathbf{u})$.

(2) (*Warm-up*) Determine whether the following parametric surfaces are regular:

(a) *Paraboloid*:

$$\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad \sigma(\mathbf{u}, \mathbf{v}) = (\mathbf{u}, \mathbf{v}, \mathbf{u}^2 + \mathbf{v}^2).$$

(b) (*Polar*) *xy-plane*:

$$\mathbf{P} : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad \mathbf{P}(\mathbf{u}, \mathbf{v}) = (\mathbf{u} \cos \mathbf{v}, \mathbf{u} \sin \mathbf{v}, 0).$$

(c) *One-sheeted hyperboloid*:

$$\mathbf{H} : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad \mathbf{H}(\mathbf{u}, \mathbf{v}) = (\cos \mathbf{u} \cosh \mathbf{v}, \sin \mathbf{u} \cosh \mathbf{v}, \sinh \mathbf{v}).$$

(3) (*Parametrise me!*) For each surface S and point $\mathbf{p} \in S$ below, give a parametrisation σ of S such that \mathbf{p} lies in the image of σ .

(a) *Plane:*

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid y = z\}, \quad \mathbf{p} = (1, -4, -4).$$

(b) *Ellipsoid:*

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + 4y^2 + 4z^2 = 4\}, \quad \mathbf{p} = (2, 0, 0).$$

(c) *Gabriel's Horn:*

$$S = \left\{ (x, y, z) \in \mathbb{R}^3 \mid x > 0, y^2 + z^2 = \frac{1}{x^2} \right\}, \quad \mathbf{p} = \left(1, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right).$$

(4) [Marked] Consider the following set:

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid xy + yz + xz = -2\}.$$

(a) Show that S is a surface.

(b) Give a parametrisation of S such that $(-1, 1, 0)$ lies in the image of S .

(c) Compute the tangent plane to S at $(-1, 1, 0)$.

(5) [Tutorial] Consider the *two-sheeted hyperboloid*:

$$\mathcal{H} = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 - z^2 = -1\}.$$

(a) Show that \mathcal{H} is a surface.

(b) Give a sketch of \mathcal{H} .

(c) Give a parametrisation of \mathcal{H} that passes through the point $(1, -1, \sqrt{3})$.

(d) Compute the tangent plane to \mathcal{H} at the point $(1, -1, \sqrt{3})$.

(6) (*Let's be self-sufficient*) For each of the following surfaces S and points $\mathbf{p} \in S$:

(i) Show that S is a surface.

(ii) Compute the tangent plane to S at \mathbf{p} .

(Unlike in Questions (4) and (5), you are not given a parametrisation of S . You will have to find your own in order to compute the tangent plane.)

(a) *Hyperbolic paraboloid:*

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid x = yz\}, \quad \mathbf{p} = (-6, 2, -3).$$

(b) *Cylinder:*

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + z^2 = 9\}, \quad \mathbf{p} = \left(-\frac{3}{\sqrt{2}}, 7, \frac{3}{\sqrt{2}}\right).$$

(7) (*Surfaces of revolution*) Let $f : (a, b) \rightarrow \mathbb{R}$ be a smooth function satisfying $f(x) > 0$ for every $x \in (a, b)$. From f , we can then define the set

$$\mathcal{R} = \{(x, y, z) \in \mathbb{R}^3 \mid a < x < b, y^2 + z^2 = [f(x)]^2\}.$$

In particular, \mathcal{R} is the *surface of revolution* obtained by taking the graph of f (in the xy -plane) and rotating it (in 3-dimensional space) around the x -axis.

(a) Show that \mathcal{R} is indeed a surface.

(b) Give a parametrisation of \mathcal{R} whose image is all of \mathcal{R} .

(c) Compute the tangent plane to \mathcal{R} at the point $(x, 0, f(x))$, for any $x \in (a, b)$.

(8) (*Fun with stereographic projections*) Consider the parametric surface

$$\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad \sigma(\mathbf{u}, \mathbf{v}) = \mathbf{p},$$

where \mathbf{p} is the (unique) point of $\mathbb{S}^2 \setminus \{(0, 0, 1)\}$ that lies on the line through the points $(\mathbf{u}, \mathbf{v}, 0)$ and $(0, 0, 1)$. (The function σ is called the *inverse stereographic projection*.)

(a) Show that σ can be described by the formula

$$\sigma(\mathbf{u}, \mathbf{v}) = \left(\frac{2\mathbf{u}}{1 + \mathbf{u}^2 + \mathbf{v}^2}, \frac{2\mathbf{v}}{1 + \mathbf{u}^2 + \mathbf{v}^2}, \frac{-1 + \mathbf{u}^2 + \mathbf{v}^2}{1 + \mathbf{u}^2 + \mathbf{v}^2} \right), \quad (\mathbf{u}, \mathbf{v}) \in \mathbb{R}^2.$$

(b) Show that σ is both injective and regular.

(c) Show that the image of σ is precisely $\mathbb{S}^2 \setminus \{(0, 0, 1)\}$.

(d) Use your knowledge of σ to construct the sphere \mathbb{S}^2 using only two regular and injective parametric surfaces.