

MTH5113 (Winter 2022): Problem Sheet 9

All coursework should be submitted individually.

- Problems marked “[**Marked**]” should be submitted and will be marked.

Be sure to write your **name** and **student ID** on your submission!

Please submit the completed problem(s) on QMPlus:

- At the portal for **Coursework Submission 5**.
-

(1) (*Warm-up*)

(a) Consider the (real-valued) function

$$F : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad F(x, y, z) = xy^2z^3,$$

as well as the parametric surface

$$\mathbf{P} : (0, 1) \times (0, 1) \rightarrow \mathbb{R}^3, \quad \mathbf{P}(u, v) = (1, u, v).$$

Compute the surface integral of F over \mathbf{P} .

(b) Consider the (real-valued) function

$$G : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad G(x, y, z) = x^2 + y^2,$$

as well as the parametric surface

$$\tau : (0, 2\pi) \times (0, 1) \rightarrow \mathbb{R}^3, \quad \tau(u, v) = (v \cos u, v \sin u, v).$$

Compute the surface integral of G over τ .

(2) (*Intro to surface integrals*) One can also define an intermediate notion of surface integration of vector fields over *parametric surfaces*. More specifically:

Definition. Let $\sigma : \mathcal{U} \rightarrow \mathbb{R}^3$ be a parametric surface, and let \mathbf{F} be a vector field that is defined on the image of γ . We then define the *surface integral* of \mathbf{F} over σ by

$$\iint_{\sigma} \mathbf{F} \cdot d\mathbf{A} = \iint_{\mathcal{U}} \{\mathbf{F}(\sigma(u, v)) \cdot [\partial_1 \sigma(u, v) \times \partial_2 \sigma(u, v)]_{\sigma(u, v)}\} du dv.$$

(a) Consider the vector field \mathbf{F} on \mathbb{R}^3 given by

$$\mathbf{F}(x, y, z) = \left(y, z^{5800} e^{x^{2000} + 46y^{1523}}, x \right)_{(x,y,z)},$$

and let \mathbf{P} be the parametric plane

$$\mathbf{P} : (0, 1) \times (0, 1) \rightarrow \mathbb{R}^3, \quad \mathbf{P}(u, v) = (1, u, v).$$

Compute the surface integral of \mathbf{F} over \mathbf{P} .

(b) Consider the vector field \mathbf{G} on \mathbb{R}^3 given by

$$\mathbf{G}(x, y, z) = (z, z, x^2 + y^2)_{(x,y,z)},$$

and let τ be the parametric torus

$$\tau : (0, 2\pi) \times (0, 1) \rightarrow \mathbb{R}^3, \quad \tau(u, v) = (v \cos u, v \sin u, v).$$

Compute the surface integral of \mathbf{G} over τ .

(c) Consider the vector field \mathbf{H} on \mathbb{R}^3 given by

$$\mathbf{H}(x, y, z) = (-x, -y, z)_{(x,y,z)},$$

and let \mathbf{q} be the (regular) parametric surface

$$\mathbf{q} : (0, 1) \times (0, 1) \rightarrow \mathbb{R}^3, \quad \mathbf{q}(u, v) = (u, v, u^2 + v^2).$$

Compute the surface integral of \mathbf{H} over \mathbf{q} .

(3) (*A Survey of Integration*) Let S denote the set

$$S = \{(u, v, u^2 - v^2) \in \mathbb{R}^3 \mid (u, v) \in (0, 1) \times (0, 1)\}.$$

(a) Show that S is a surface. In addition, give an injective parametrisation of S whose image is precisely all of S .

(b) Compute the surface integral over S of the real-valued function

$$F : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad F(x, y, z) = xy.$$

(The double integral you get from expanding the surface integral is not so pleasant; you will probably have to use the method of substitution twice to compute it.)

- (c) Let us also assign to S the *upward-facing orientation*, i.e. the orientation in the *positive z-direction*. Then, compute the surface integral over S of the vector field

$$\mathbf{G}(x, y, z) = (xy^2, yx^2, 1)_{(x,y,z)}, \quad (x, y, z) \in \mathbb{R}^3.$$

- (4) [Marked] Let S denote the following surface:

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid x = y^2 - z^2, 0 < y < 1, 0 < z < 1\}.$$

- (a) Compute the surface integral over S of the function

$$G : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad G(x, y, z) = yz.$$

- (b) Let us also assign to S the *orientation in direction of increasing x-value*. Then, compute the surface integral over S of the vector field \mathbf{H} on \mathbb{R}^3 given by

$$\mathbf{H}(x, y, z) = (y, 0, z)_{(x,y,z)},$$

- (5) [Tutorial]

- (a) Consider the surface (*you may assume this is indeed a surface*)

$$\mathcal{P} = \{(u, v, u^4 + v) \in \mathbb{R}^3 \mid (u, v) \in (0, 1) \times (-1, 1)\}.$$

Compute the surface integral over \mathcal{P} of the following function:

$$F : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad F(x, y, z) = 6x^5.$$

- (b) Consider the sphere,

$$\mathbb{S}^2 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\},$$

and let \mathbb{S}^2 be given the “outward-facing” orientation. Compute the surface integral

over \mathbb{S}^2 of the vector field \mathbf{F} on \mathbb{R}^3 defined by the formula

$$\mathbf{F}(\mathbf{x}, \mathbf{y}, z) = (0, 0, z^3)_{(\mathbf{x}, \mathbf{y}, z)}.$$

(6) (*A-levels, revisited*)

(a) Show that the surface area of a *sphere of radius* $r > 0$,

$$S_r = \{(\mathbf{x}, \mathbf{y}, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = r^2\},$$

is equal to $4\pi r^2$.

(b) Show that the area of the side of a cone with base radius $r > 0$ and height $h > 0$,

$$C_{r,h} = \left\{ (\mathbf{x}, \mathbf{y}, z) \in \mathbb{R}^3 \mid 0 < z < h, x^2 + y^2 = r^2 \left(1 - \frac{z}{h}\right)^2 \right\},$$

is equal to $\pi r \sqrt{r^2 + h^2}$.

(7) (*Reversal of orientations*) Let $S \subseteq \mathbb{R}^3$ be an oriented surface, and let $\sigma : \mathcal{U} \rightarrow S$ be a parametrisation of S . Moreover, define the set

$$\mathcal{U}_r = \{(\mathbf{v}, \mathbf{u}) \mid (\mathbf{u}, \mathbf{v}) \in \mathcal{U}\}$$

and define the parametric surface

$$\sigma_r : \mathcal{U}_r \rightarrow \mathbb{R}^3, \quad \sigma_r(\mathbf{v}, \mathbf{u}) = \sigma(\mathbf{u}, \mathbf{v}).$$

In other words, σ_r is precisely σ but with the roles of \mathbf{u} and \mathbf{v} reversed.

(a) Show that for any $(\mathbf{u}, \mathbf{v}) \in \mathcal{U}$,

$$\partial_1 \sigma_r(\mathbf{v}, \mathbf{u}) \times \partial_2 \sigma_r(\mathbf{v}, \mathbf{u}) = -[\partial_1 \sigma(\mathbf{u}, \mathbf{v}) \times \partial_2 \sigma(\mathbf{u}, \mathbf{v})].$$

(b) Show that σ_r is also a parametrisation of S , and that σ_r has the same image as σ .

(c) Use the formula from part (a) to conclude that if σ generates an orientation \mathbf{O} of S , then σ_r generates the orientation opposite to \mathbf{O} .

(8) (*The paradox of Gabriel's horn*) Consider the surface of revolution

$$\mathbf{G} = \left\{ (x, y, z) \in \mathbb{R}^3 \left| y^2 + z^2 = \frac{1}{x^2}, x > 1 \right. \right\},$$

which is sometimes nicknamed *Gabriel's horn*. (*Before proceeding, you should search for "Gabriel's horn" on Google Images to see an illustration of \mathbf{G} .*)

(a) Show that \mathbf{G} has infinite surface area.

(b) Show that the interior of \mathbf{G} ,

$$\mathbf{I} = \left\{ (x, y, z) \in \mathbb{R}^3 \left| y^2 + z^2 \leq \frac{1}{x^2}, x > 1 \right. \right\},$$

has finite volume.

In other words, *you can fill up the inside of the "horn" with a finite amount of paint, but you cannot paint the "horn" itself using a finite amount of paint!*