

MTH5113 (Winter 2022): Problem Sheet 10

All coursework should be submitted individually.

- Problems marked “[**Marked**]” should be submitted and will be marked.

*Be sure to write your **name** and **student ID** on your submission!*

Please submit the completed problem(s) on QMPlus:

- At the portal for **Coursework Submission 5**.
-

(1) (*Warm-up*) Consider the following constrained optimisation problem:

- *Maximise the function $f(x, y) = x$, subject to the constraint $x^2 + y^2 = 1$.*
- (a) Give the solution to the above problem *without doing any calculations. (The answer should be obvious from inspection alone; draw a picture if you are not sure.)*
- (b) Solve the above problem using the method of Lagrange multipliers. Verify that your solution matches what you deduced in part (a).

(2) (*Warm-up*) Consider the following constrained optimisation problem:

- *Minimise the function $f(x, y, z) = z$, subject to the constraint $x^2 + y^2 + z^2 = 1$.*
- (a) Give the solution to the above problem *without doing any calculations. (The answer should be obvious from inspection alone; draw a picture if you are not sure.)*
- (b) Solve the above problem using the method of Lagrange multipliers. Verify that your solution matches what you deduced in part (a).

(3) [**Marked**] Solve the following problem using the method of Lagrange multipliers:

- *Find the maximum and minimum values of $x - y^3$, subject to the constraint $x^2 + 9y^2 = 9$. At which points are the maximum and minimum values achieved?*

(4) (*Differential Geometry and Game Theory*) Let x^2 be the number of hours of *MTH5113* lectures and tutorials you attend, and let y^2 be the number of hours of *MTH5113* lectures and tutorials you skip. As you know, the *constraint* is that there are only 43 total hours of lectures and tutorials in *MTH5113*. Now, suppose that the “effectiveness” of your learning in *MTH5113*, as a function of the above hours spent, is modelled by the relation

$$E = 100x^2 + y^2.$$

(Here, a higher value of E means better learning!) Your objective here is to *optimise* the “effectiveness” of your learning experience in *MTH5113*!

- (a) Express the above objective as a constrained optimisation problem.
- (b) Use the method of Lagrange multipliers to solve the problem in part (a).
- (c) Given your answer in (b), what optimal strategy should you adopt in order to have the most effective learning experience in *MTH5113*? :)

(5) [Tutorial] Use the method of Lagrange multipliers to solve the following:

- (a) Find the minimum and maximum of $4x^2 - y^2$, subject to the constraint $x^2 + 4y^2 = 4$.
- (b) Find the unit vectors $(x, y, z) \in \mathbb{R}^3$ that maximise and minimise the dot product,

$$(6, -3, 2) \cdot (x, y, z).$$

(6) (*Conservative and liberal vector fields*)

- (a) Let f be the real-valued function

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad f(x, y, z) = x^4 y^2 z.$$

Compute the integral of the vector field ∇f over the curve

$$C = \{(t, t^2, t^3) \in \mathbb{R}^2 \mid t \in (0, 1)\},$$

where C is given the *rightward* (i.e. in the direction of increasing x -value) *orientation*.

- (b) Let $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ be given by the formula

$$g(x, y) = x^{17} e^{y+x^2 y^5 \cos x^7} + y^4 + e^{x^2+y^2+x^{42}+y^{1776}} e^{y x^2}.$$

Find the integral of the vector field ∇g over the unit circle

$$\mathcal{C} = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^2 \mid \mathbf{x}^2 + \mathbf{y}^2 = 1\},$$

where $\mathcal{C}$ is given the *anticlockwise orientation*.

(c) Integrate the vector field

$$\mathbf{H}(\mathbf{x}, \mathbf{y}) = (-\mathbf{y}, \mathbf{x}), \quad (\mathbf{x}, \mathbf{y}) \in \mathbb{R}^2,$$

over the unit circle $\mathcal{C}$ from part (b), where $\mathcal{C}$ again has the anticlockwise orientation.

(d) From your answer in part (c), conclude that the vector field $\mathbf{H}$ cannot be the gradient of any real-valued function $h : \mathbb{R}^2 \rightarrow \mathbb{R}$.

(7) (*Lagrangian Formulation of Multipliers*) Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ be smooth functions, and fix $c \in \mathbb{R}$. Show that the following conditions are equivalent:

(i) $(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^2$ and $\lambda \in \mathbb{R}$ satisfy the following system of equations:

$$\nabla f(\mathbf{x}, \mathbf{y}) = \lambda \cdot \nabla g(\mathbf{x}, \mathbf{y}), \quad g(\mathbf{x}, \mathbf{y}) = c.$$

(ii) $(\mathbf{x}, \mathbf{y}, \lambda) \in \mathbb{R}^3$ satisfies the equation

$$\nabla \mathcal{L}(\mathbf{x}, \mathbf{y}, \lambda) = (0, 0, 0)_{(\mathbf{x}, \mathbf{y}, \lambda)},$$

where the function $\mathcal{L}$, called the *Lagrangian*, is defined by

$$\mathcal{L} : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad \mathcal{L}(\mathbf{u}, \mathbf{v}, w) = f(\mathbf{u}, \mathbf{v}) - w[g(\mathbf{u}, \mathbf{v}) - c].$$

(Thus, the method of Lagrange multipliers could also be formulated in terms of $\mathcal{L}$ in (ii)—if the maximum or minimum of f , subject to the constraint g , is achieved at $(\mathbf{x}, \mathbf{y})$, then there is some $\lambda \in \mathbb{R}$ such that $(\mathbf{x}, \mathbf{y}, \lambda)$ is a critical point of the Lagrangian $\mathcal{L}$.)

(8) (*Multiple Constraints*) Assume the following formal setting:

- Let $\mathbf{U} \subseteq \mathbb{R}^3$ be open and connected.
- Let $f : \mathbf{U} \rightarrow \mathbb{R}$, $g : \mathbf{U} \rightarrow \mathbb{R}$, $h : \mathbf{U} \rightarrow \mathbb{R}$ be smooth functions.

- Suppose $\nabla g(\mathbf{p}) \times \nabla h(\mathbf{p})$ is nonvanishing at every $\mathbf{p} \in \mathcal{U}$.

Under the above assumptions, the following result holds:

- **Theorem.** Suppose f achieves its maximum or minimum value on

$$\mathcal{C} = \{(\mathbf{x}, \mathbf{y}, \mathbf{z}) \in \mathcal{U} \mid g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \mathbf{c}, h(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \mathbf{d}\}$$

at a point $\mathbf{p} \in \mathcal{C}$. Then, there exist $\lambda, \mu \in \mathbb{R}$ such that

$$\nabla f(\mathbf{p}) = \lambda \cdot \nabla g(\mathbf{p}) + \mu \cdot \nabla h(\mathbf{p}).$$

Using the preceding theorem:

- Devise a corresponding *method of Lagrange multipliers* for solving the following constraint optimisation problem: *maximise or minimise* $f(\mathbf{x}, \mathbf{y}, \mathbf{z})$, *subject to the simultaneous constraints* $g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \mathbf{c}$ *and* $h(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \mathbf{d}$.
- Using your strategy from part (a), find the maximum and minimum values of $\mathbf{x} + \mathbf{y} + \mathbf{z}$, subject to the simultaneous constraints $\mathbf{x}^2 + \mathbf{y}^2 = 1$ and $\mathbf{x} - \mathbf{z} = 1$.

(>9000) (*Extra Exploration*) Put your geometry, calculus, and linear algebra knowledge to the test! Can you prove the theorem stated in Question (8)?

(Hint: The starting point is to observe that $\mathcal{C}$ is a curve, by the result of Question (9) of Problem Sheet 4. From here, you have all the background you need to do this!)

(Note: While I will not be posting the solution to this problem, I would be happy to chat with anyone who wishes to attempt it. Good luck!)