

## MTH5113 (Winter 2022): Problem Sheet 11

These problems are for practice only. *You do not need to submit them.*

(1) (*Boring computations—but you should know how to do them*) Consider the following vector fields  $\mathbf{F}$ ,  $\mathbf{G}$ ,  $\mathbf{H}$  on  $\mathbb{R}^3$ , where:

$$\mathbf{F}(x, y, z) = (x, y, z)_{(x,y,z)},$$

$$\mathbf{G}(x, y, z) = (x^2, -2xy, 3xz)_{(x,y,z)},$$

$$\mathbf{H}(x, y, z) = (x^2 + y^2 + z^2, x^4 - y^2 z^2, xyz)_{(x, y, z)}.$$

- (a) Compute the divergence of  $\mathbf{F}$ ,  $\mathbf{G}$ , and  $\mathbf{H}$  at each point  $(x, y, z) \in \mathbb{R}^3$ .
- (b) Compute the curl of  $\mathbf{F}$ ,  $\mathbf{G}$ , and  $\mathbf{H}$  at each point  $(x, y, z) \in \mathbb{R}^3$ .

(2) (*Fun with Green's theorem*) Let  $\mathbb{C}$  denote the circle

$$C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 9\},$$

and let us assign to  $\mathbb{C}$  the anticlockwise orientation.

- (a) Let  $\mathbf{F}$  be the vector field on  $\mathbb{R}^2$  defined by

$$\mathbf{F}(x, y) = (x - y, x + y)_{(x, y)}.$$

Compute the curve integral of  $\mathbf{F}$  over  $C$  *directly*.

- (b) Compute the curve integral from part (a) *using Green's theorem*. Check that your answer here matches the answer you obtained in part (a).
- (c) Let  $\mathbf{G}$  be the vector field on  $\mathbb{R}^2$  defined by

$$\mathbf{G}(\mathbf{x}, \mathbf{y}) = (\mathbf{x}^{9999999999999999} e^{\mathbf{x}} + \mathbf{y}, \mathbf{x} + \mathbf{y}^{3333333333333333} e^{2\mathbf{y}})_{(\mathbf{x}, \mathbf{y})}.$$

Compute (using your favourite method) the curve integral of  $\mathbf{G}$  over  $C$ .

**(3)** (*Fun with Stokes' theorem*) Let  $S$  denote the upper half-sphere,

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1, z > 0\}.$$

In addition, let  $C$  denote the boundary of  $S$ , i.e. the circle

$$C = \{(x, y, 0) \in \mathbb{R}^3 \mid x^2 + y^2 = 1\},$$

and assign to  $C$  the (anticlockwise) orientation generated by the parametrisation

$$\gamma : \mathbb{R} \rightarrow \mathbb{R}^3, \quad \gamma(t) = (\cos t, \sin t, 0).$$

(a) Let  $\mathbf{F}$  be the vector field on  $\mathbb{R}^3$  given by

$$\mathbf{F}(x, y, z) = (x, y, z)_{(x,y,z)}.$$

Compute directly the curve integral of  $\mathbf{F}$  over  $C$ .

(b) Evaluate the integral from part (a) by applying *Stokes' theorem* and computing instead an appropriate integral over  $S$ . Make sure that you obtain the same answer as in (a).

(c) Let  $\mathbf{G}$  and  $\mathbf{H}$  be smooth vector fields on  $\mathbb{R}^3$ , and assume that  $\mathbf{G}(\mathbf{p}) = \mathbf{H}(\mathbf{p})$  for every  $\mathbf{p} \in C$ . Using Stokes' theorem, show that

$$\iint_S (\nabla \times \mathbf{G}) \cdot d\mathbf{A} = \iint_S (\nabla \times \mathbf{H}) \cdot d\mathbf{A}.$$

(Here,  $S$  can be assigned either of its orientations.)

(4) (*Fun with the divergence theorem*) Let  $\mathbb{S}^2$  denote the unit sphere centred at the origin,

$$\mathbb{S}^2 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\},$$

and assign to  $\mathbb{S}^2$  the outward-facing orientation.

(a) Let  $\mathbf{F}$  be the vector field on  $\mathbb{R}^3$  given by

$$\mathbf{F}(x, y, z) = (x, y, z)_{(x,y,z)}.$$

Compute directly the surface integral of  $\mathbf{F}$  over  $\mathbb{S}^2$ .

(b) Evaluate the integral from part (a) by applying the *divergence theorem* and computing an appropriate triple integral. Make sure that you obtain the same answer as in (a).

(c) Let  $\mathbf{L}$  be the vector field on  $\mathbb{R}^3$  given by

$$\mathbf{L}(x, y, z) = \left( y^{543} e^{y^2+z^4} z^{5234}, e^{x^{562} z^{27} - x^{12} z^{43}} (x + z e^x)^{127}, 1 + x^{10} y + 24 y^{17} e^{42 y^3} \right)_{(x,y,z)}.$$

Evaluate the surface integral of  $\mathbf{L}$  over  $\mathbb{S}^2$ .

(5) (*Setting boundaries I*)

(a) Describe the boundaries of the following subsets of  $\mathbb{R}^2$ , as one or more curves in  $\mathbb{R}^2$ :

- (i)  $D_1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + 4y^2 < 4\}$ .
- (ii)  $D_2 = (0, 1) \times (0, 2)$ .
- (iii)  $D_3 = \{(x, y) \in \mathbb{R}^2 \mid x < 1, y > 0, y < x\}$ .

(b) Describe the boundaries of the following surfaces in  $\mathbb{R}^3$ , as one or more curves in  $\mathbb{R}^3$ :

- (i)  $S_1 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1, x > 0\}$ .
- (ii)  $S_2 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 = 1, -1 < z < 1\}$ .
- (iii)  $S_3 = \{(x, y, 0) \in \mathbb{R}^3 \mid x \in (0, 1), y \in (0, 2)\}$ .

(c) Describe the boundaries of the following regions in  $\mathbb{R}^3$ , as one or more surfaces in  $\mathbb{R}^3$ :

- (i)  $V_1 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 < 9\}$ .
- (ii)  $V_2 = (0, 1) \times (0, 1) \times (0, 1)$ .
- (iii)  $V_3 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 < 1, -1 < z < 1\}$ .

(6) (*Setting boundaries II*)

(a) Suppose you apply Green's theorem to each of the regions  $D_i$  in Question (5a). Describe the resulting orientation(s) on the boundary of each  $D_i$  that you would obtain, according to the statement of Green's theorem.

(b) Suppose you apply Stokes' theorem to each of the surfaces  $S_i$  in Question (5b), where:

- For  $S_1$ , we use the orientation facing the positive  $x$ -direction.
- For  $S_2$ , we use the outward-facing orientation.
- For  $S_3$ , we use the orientation facing the positive  $z$ -direction.

Describe the resulting orientation(s) on the boundary of each  $S_i$  that you would obtain, according to the statement of Stokes' theorem.

- (c) Suppose you apply the divergence theorem to each of the regions  $V_i$  in Question (5c). Describe the resulting orientation(s) on the boundary of each  $V_i$  that you would obtain, according to the statement of the divergence theorem.

(7) (*Second derivative identities*)

- (a) Let  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  be a smooth function. Show that the curl of the gradient of  $f$  vanishes everywhere, that is, show that for any  $\mathbf{p} \in \mathbb{R}^3$ ,

$$[\nabla \times (\nabla f)](\mathbf{p}) = (0, 0, 0)_{\mathbf{p}}.$$

- (b) Let  $\mathbf{F}$  be a smooth vector field on  $\mathbb{R}^3$ . Show that the divergence of the curl of  $\mathbf{F}$  vanishes everywhere, that is, show that for any  $\mathbf{p} \in \mathbb{R}^3$ ,

$$[\nabla \cdot (\nabla \times \mathbf{F})](\mathbf{p}) = 0.$$

(8) (*Connections to Complex Variables*) (*Not examinable*) Consider the plane  $\mathbb{R}^2$ , or equivalently, the complex plane  $\mathbb{C}$ . Let  $C$  denote the unit circle centred at the origin,

$$C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\} \simeq \{z \in \mathbb{C} \mid |z| = 1\},$$

and assign to  $C$  the anticlockwise parametrisation. Furthermore, let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a complex-analytic function, and write  $f$  in terms of its components as

$$f(z) = u(z) + i \cdot v(z), \quad z \in \mathbb{C}.$$

- (a) Express the real and imaginary parts of the contour integral of  $f$  over  $C$ ,

$$\operatorname{Re} \int_C f(z) \, dz, \quad \operatorname{Im} \int_C f(z) \, dz,$$

as curve integrals of appropriate vector fields over  $C$ .

- (b) Use Green's theorem to prove Cauchy's theorem on  $C$ :

$$\int_C f(z) \, dz = 0.$$

(Hint: Recall that  $u$  and  $v$  satisfy the Cauchy-Riemann equations.)

(9) (Green's theorem fail) Let  $C$  be as in Question (8), and let  $\mathbf{F}$  be the vector field

$$\mathbf{F}(x, y) = \left( -\frac{y}{x^2 + y^2}, \frac{x}{x^2 + y^2} \right)_{(x,y)}, \quad (x, y) \in \mathbb{R}^2 \setminus \{(0, 0)\}.$$

(a) Show that for any  $(x, y) \in \mathbb{R}^2 \setminus \{(0, 0)\}$ ,

$$\partial_x \left( \frac{x}{x^2 + y^2} \right) - \partial_y \left( -\frac{y}{x^2 + y^2} \right) = 0.$$

(b) On the other hand, show that the integral of  $\mathbf{F}$  over  $C$  is *not* zero. Why does this not contradict the statement of Green's theorem? (More specifically, why do Green's theorem and part (a) not imply that the integral of  $\mathbf{F}$  over  $C$  is zero?)

(c) (Not examinable) For those taking *Complex Variables*, can you relate what you saw in parts (a) and (b) to some contour integrals that you have seen?